

Review of Gray's beaked whale (*Mesoplodon grayi*) in the South East Pacific

JHOANN CANTO HERNÁNDEZ^{1,2*}, GUILLERMO D'ELIA³, RICHARD CADENILLAS^{3,4}, FREDERICK TORO⁵, CINTYA BORRONI⁴, GERARDO CERDA⁵, AND SONIA ESPAÑOL-JIMÉNEZ⁶

¹Área Zoología Vertebrados, Museo Nacional de Historia Natural, Casilla 787, Santiago, Chile. E-mail: jhoann.canto@mnhn.gob.cl (JCH).

²Medicina Veterinaria, Universidad Santo Tomás, Santiago, Chile.

³Instituto de Ciencias Ambientales y Evolutivas, Universidad Austral de Chile, Valdivia, Chile. E-mail: guille.delia@gmail.com (GD); cadenillasordinola@gmail.com (RC).

⁴Universidad Señor de Sipán, Chiclayo - Perú. E-mail: cadenillasordinola@gmail.com (RC).

⁵Facultad de Medicina Veterinaria y Recursos Naturales. Universidad Santo Tomás, Viña del Mar, Chile. E-mail: Frederick.Toro.C@gmail.com (FT).

⁶Escuela de Medicina Veterinaria. Facultad de Ciencias de la Vida, Universidad Andres Bello, Chile. E-mail: cintya.borroni@unab.cl (CB).

⁷Servicio Nacional de Pesca (SERNAPESCA), Videla 240 Piso 6, Región Coquimbo, Chile. E-mail: gcerda@sernapesca.cl (GC).

⁸Melimoyu Ecosystem Research Institute. Av. Kennedy 5682, Santiago, Chile. E-mail: sespanol@fundacionmeri.cl (SE-J).

*Corresponding author

Information collected from a complete female juvenile individual of Gray's beaked whale (*Mesoplodon grayi*) stranded on the Guanaqueros coast, Coquimbo Region in Chile (30°S) is provided. Difficulties to differentiate specimens of Gray's beaked whale and Hector's beaked whale (*M. hectori*) are discussed based on the use of diagnostic phenotypic characters, such as differences in color patterns and position of teeth on the lower jaw. The identification of the studied specimen as Gray's beaked whale was supported by a detailed review of cranial characters and molecular analyses. Finally, we provide an updated list containing all known Chilean records for this species.

Se proporciona información del registro más completo de *Mesoplodon grayi* (Ziphiidae) que se conoce para Chile y que corresponde a una hembra juvenil varada en la costa de Guanaqueros, Región de Coquimbo (30°S). Se discuten los problemas de determinación de *Mesoplodon grayi* con respecto a *M. hectori* en base a la utilización de caracteres fenotípicos generales como patrones de coloración. La asignación a *Mesoplodon grayi* se apoyó en la revisión detallada de caracteres craneales y un análisis molecular. Además, se presenta un listado actualizado con todos los registros conocidos para esta especie en las costas de Chile.

Keywords: Distribution; Gray's beaked whale; Odontoceti; South America; Ziphiidae.

© 2022 Asociación Mexicana de Mastozoología, www.mastozoologiamexicana.org

Introduction

Strandings represent a unique opportunity to collect information about elusive cetaceans (Canto *et al.* 1991; Leeney *et al.* 2008; Pikesley *et al.* 2011). Although these events are frequent across different coastal areas of the planet (Evans *et al.* 2005; Berta *et al.* 2015; Moore *et al.* 2018), the frequency of such stranding events differs among species (Leeney *et al.* 2008). In the case of species of the family Ziphiidae, records are scarce, and many of them are based on the stranding of specimens in an advanced state of decay or isolated bone remains (Litcher 1986; Dalebout *et al.* 1988). Therefore, only a few high-quality photographic records of ziphiids exist, given that instances of complete individuals in good condition are exceptional (Cox *et al.* 2006). Traditionally, identifying specimens of *Mesoplodon* at the species level has been a complex task (Dalebout *et al.* 2002, 2004; Van Helden *et al.* 2002; Cappozzo *et al.* 2005). For example, relying on body patterns of coloration (a trait usually used for species identification) can sometimes result in wrong taxonomic identifications (Helden *et al.* 2002; Dalebout *et al.* 2014) due to the insufficient knowledge about intra and interspecific color variation as well as a poor understanding of how coloration varies after death.

The ziphiid genus *Mesoplodon* is the most diverse, accounting for 15 of the 22 species of the family (Cappozzo *et al.* 2005; McLeod *et al.* 2006; Pitman 2009; McLeod 2009, 2017; Yamada *et al.* 2019). Ten species of *Mesoplodon* are known from the Atlantic and Pacific oceans of South America; these are *M. bowdoini* (Andrews' beaked whale), *M. densirostris* (Blainville's beaked whale), *M. europaeus* (Gervais' beaked whale), *M. ginkgodens* (Ginkgo-toothed beaked whale), *M. grayi* (Gray's beaked whale), *M. hectori* (Hector's beaked whale), *M. layardii* (Strap-toothed beaked whale), *M. mirus* (True's beaked whale), *M. peruvianus* (Pygmy beaked whale), and *M. traversii* (Spade-toothed whale; McLeod *et al.* 2006). Six species of *Mesoplodon* have been reported in Chile: *M. densirostris*, *M. grayi*, *M. hectori*, *M. layardii*, *M. peruvianus*, and *M. traversi* (Canto and Yáñez 2009; D'Elia *et al.* 2020). Reports of these species in the Chilean coast are based on less than 20 records, a fact that highlights the importance of documenting any newly emerging record for this genus. These records are based mainly on bone remains (Sielfeld 1979, 1980, 1983; Cárdenas *et al.* 1986; Canto *et al.* 1992; Reyes *et al.* 1995; Sanino *et al.* 2007) and in only a few cases, on specimens in good condition. Among the latter, there is a male of Strap-toothed beaked whale

stranded at Caleta Tortel, Aysén Region in 2019 (personal communication S.E-J.), in addition to the new specimen of *M. grayi* we are reporting in this work.

Information on Gray's beaked whales consists of 192 records held in GBIF (2019) worldwide. A single record, not included in GBIF, is known from the South American Pacific Ocean, from the Peruvian coast (Reyes 1990). In Chile, Mann (1958) initially reported this species. Later, Venegas and Sielfeld (1978) and Sielfeld (1979) documented for the first time the presence of this species in the country, based on osteological material found at the Strait of Magellan. So far, these are the only records of this species for Chile concentrated in the southern part of the country.

Here we report a new Chilean record for Gray's beaked whales on the basis of a specimen initially identified as *M. hectori* by the Servicio Nacional de Pesca in 2018. This new record, from Guanaqueros beach (Figure 1), confirms the wide distribution of the species in the South-East Pacific, as suggested by MacLeod *et al.* (2006). The first DNA sequence of any Chilean specimen of this species is also provided. The analysis of skull morphology and coloration suggests that the species presents different morphs, which can be attributed to sexual dimorphism or geographic variation.

Materials and methods

On December 21, 2018, a young female odontocete stranded in the coast of Guanaqueros, Coquimbo Region, Chile, was collected by SERNAPESCA and later transferred to the Museo Nacional de Historia Natural (MNHN), Santiago. A necropsy was performed to determine the possible causes of death. External measurements were taken according to Reyes and Waerebeek (2018), while cranial measurements were taken following Ross (1984). Muscle tissue samples were fixed in 95 % ethanol. Fat samples, the respiratory system, the heart, the uterus and the ovaries were frozen at -18° C. All bone material, as well as the samples, are housed in the Mammal Collection of the MNHN under catalogue number MNHN/MAM 1972.

For the specific determination of specimen MNHN/MAM 1972, comparisons were made with the descriptions and morphometric data of adult specimens of *M. hectori*, *M. peruvianus*, *M. densirostris*, *M. bowdoini*, *M. perrini*, *M. stejnegeri*, *M. grayi*, and *M. europaeus* (Ross 1970; Venegas and Sielfeld 1978; Sielfeld 1979; 1983; Lichter 1986; Mead and Baker 1987; Nobuyuki *et al.* 1987; Mead 1989; Baker and Helden 1999; Norman and Mead 2001; Zerbini and Secchi 2001; Dalebout *et al.* 2002; Capazzo *et al.* 2005; Laporta *et al.* 2005).

In addition, genetic analyses were performed based on a fragment (801 bp) of the mitochondrial gene encoding the cytochrome b (cytb). Total genomic DNA was extracted from a sample of fresh muscle tissue using the Wizard SV Genomic DNA Purification System (Promega ©). The cytb fragment was amplified using primers MVZ 05 and MVZ 16 following the protocol mentioned in Chiquito *et al.* (2014).




Figure 1. Map showing the three confirmed records of *Mesoplodon grayi* from the Chilean coast. The unconfirmed record from Chanavayita in northern Chile is also signalled. The record presented in this contribution corresponds to that of Guanaqueros.

The new DNA sequence was edited, including checking for the existence of internal stop codons and changes in the reading frame, using CodonCode (Codon-Code, Dedham, MA, USA). The new DNA sequences were deposited in GenBank under accession number MT813067.

Sequence alignment was done with MAFFT v.7.310 (Kato and Standley 2013) using L-INS-i as an iterative refinement method (Kato and Toh 2008). Model TIM2+F+I+G4 was selected with IQ-TREE v.1.6.5 (Trifinopoulos *et al.* 2016). Two phylogenetic methods were used. Maximum Likelihood (ML) analysis was conducted in IQ-TREE with perturbation strength was set to 0.5 and the number of unsuccessful iterations set to 100; nodal support was evaluated by 1,000 ultrafast Bootstrap replicates (BS; Akaike 1973). A Bayesian analysis (BI) was conducted in MrBayes 3.2 (Ronquist and Huelsenbeck 2013) by means of two independent runs, each with five heated and one cold Markov chains and all model parameters estimated in MrBayes. Uniform-interval priors were assumed for all parameters except base composition and substitution model parameters, which considered a Dirichlet prior. Runs lasted 50 million generations, with trees sampled every 1,000 generations. Convergence was checked by plotting log-likelihood values against generation time. The first 25 % of the trees generated were discarded as burned, and only the remaining trees were used to compute a 50 % majority rule consensus tree and to obtain posterior probability (PP) estimates for each clade. Both analyses included sequences of all members of the family Ziphiidae, and *Physeter catodon* was employed as an outgroup; see supplementary material 1 for the accession numbers.

A list of Chilean strandings and other records involving species of *Mesoplodon* was compiled from the literature and qualified reports. References to unsupported sightings were excluded.

Results

According to SERNAPESCA, specimen MNHN/MAM 1972 would have died very shortly before stranding. This statement is based on its general good conditions, including coloration and absence of external scars or bruises (Figure 2A-B). The necropsy revealed the skull was severely damaged (see supplementary material 2).

Specimen MNHN/MAM 1972 corresponds to a juvenile female, as cranial sutures are not completely fused, especially the front-supraoccipital one. Furthermore, the skull presents a globose shape at the parietal level, which is characteristic of juvenile cetaceans (Del Castillo 2016). In the nasal region, at the vertex level, a posteroanterior projection of smaller size is observed. The mesorostral canal is cylindrical and does not present ossification like adult specimens of the same species and other species of the genus (Besharse 1971).

Body and skull-mandibular measurements of specimen MNHN/MAM1972 are presented in Table 1. The specimen measured 2.47 m and weighted 185 k. The body has a spindle shape, showing the largest circumference around its midpoint. The beak is short and narrow with the line of the mouth curved shaped (Figure 2B). There is a black patch around the eyes that projects towards the top of the head, in

front of the respiratory opening, but with a less marked coloration, tending to grey. The ventral zone of the mandibular (except the area of the beak) and the gular region is greyish-white. The beak is a dark brown that fades slightly towards the melon area where it intersects with grey tones. Behind the respiratory opening, it presents a light pale cream color with a variation of shades in the form of irregular patches. The dorsal area of the body also presents a light beach color with variation in the shades that extend along the entire back. Ventrally, the coloration is greyish white and extends to the caudal peduncle. The dorsal fin is dark grey. The dorsal surface of the pectoral fins has grey color with a variation of shades to darker tones. The caudal lobes in dorsal view are grey, whilst on the ventral side, are light grey.

In the skull the anterior border of the pterygoid sinus reaches the level of the maxillary foramina (= dorsal infra-orbital foramina according to Mead and Fordyce 2009). The left premaxillary ridge is not marked and is smaller than the one in the right (Figure 2C, D, E and F). A pair of teeth located in the area of the mandibular symphysis is evident. The jaw is very short compared to adult specimens of the same species (Figure 2G-H). Both the mandibular condyle and coronoid process were not preserved in both mandibular branches.

The ML and BI analyses showed that the relationship between the members of the family Ziphiidae are mostly unresolved and poorly supported; the genus *Mesoplodon* was recovered as paraphyletic with respect to *Indopacetus pacificus*. However, all the species of *Mesoplodon* were recov-




Figure 2. A) View of specimen MNHN / MAM 1972 at its site of stranding on Guanaqueros beach, Coquimbo Region, Chile. The literature mentions a dark brown, dark grey or black colouration pattern on the dorsal surface and paler on the ventral surface, with the tip of the bill being generally white, unlike the rest of the body (Wilson and Mittermeir 2014). Specimen MNHN/MAM 1972 shows a different colour pattern with a dark-coloured peak (see black arrow). The preservation of the colouration stands out because photo was taken only a few hours after the specimen stranded. B) Detail of the head of specimen MNHN / MAM 1972 showing its general colouration and the shape of the beak. C-F) Dorsal and ventral view of the skull of (C and E) an adult female specimen of *Mesoplodon grayi* (Smithsonian Museum of Natural History Mammal Collection USNM A 49880) and (D and F) specimen MNHN/MAM 1972. G-H) Dorsal view of the mandible of (G) an adult female *Mesoplodon grayi* (USNM A 49880) and (H) specimen MNHN/MAM 1972, without the mandibular condyle and coronoid process (black arrows). In both specimens, it is observed that the position of the teeth is immediately in front of the mandibular symphysis (red arrows).

ered monophyletic and with high support. The sequence obtained for MNHN/MAM 1972 falls in the clade formed by sequences of *M. grayi* (BS = 99; PP = 1; Figure 3). The sequence obtained from MNHN/MAM 1972 diverges on average from other sequences of *Gray's beaked whales* by 1.45 %, while divergence with that of *M. perrini* is 8.97 % and *M. peruvianus* 10.71 %. In the sequence distance matrix by species, values ranging between 0.00783 and 0.0225 are observed for four *M. grayi* sequences (see supplementary material 4). *M. grayi* form a clade (BS = 95; PP=0.89) with *M. peruvianus* and *M. perrini*.

Twenty-two records of the genus *Mesoplodon* have been reported for Chile, only three of which have been identified as *M. grayi*, and additionally, a record is indicated on the coast of Iquique for the species (Figure 1, Table 1).

Discussion

Chilean records of *Mesoplodon*, based on sightings or on strandings, are scarce. Compared to other cetaceans, such as *Pseudorca*, *Tursiops*, *Lagenorhynchus*, *Cephalorhynchus*, and *Phocoena*, these records are considerably less frequent (Aguayo 1975). Information compiled by SERNAPESCA (2021) between 2009 and 2021 include 1.093 strandings of cetaceans in Chile, of which only two correspond to species of Ziphiidae. One of the main causes of the low frequency of occurrences is that species of *Mesoplodon*, and more generally of ziphiids, mainly inhabit the oceanic zone, reaching only occasionally the continental coasts. However, during 2018 and 2020 both sightings and strandings of various species of this family have increased throughout

Table 1. Records of species of the genus *Mesoplodon* along the Chilean coast (1968-2020).

Specie	Locality	Date	Reference	Comment
<i>M. hectori</i>	Windhond bay, Navarino island, Region of Magallanes and Chilean Antarctica	March, 1979	Sielfeld 1979	Skull housed in Zoological collection Instituto de la Patagonia (CE-24)
	Windhond bay, Navarino island, Region of Magallanes and Chilean Antarctica	March, 1979	Sielfeld 1979	Skull housed in Zoological collection Instituto de la Patagonia (CE-25)
<i>M. layardii</i>	Rio Seco, Punta Arenas, Region of Magallanes and Chilean Antarctica	1968	Venegas and Sielfeld 1978	Picture of specimen housed in files of Instituto de la Patagonia, Universidad de Magallanes
	Cabo Espiritu Santo, Tierra del Fuego	February 15, 1978	Venegas and Sielfeld 1978	Right tooth housed in Zoological collection Instituto de la Patagonia Universidad de Magallanes (CE-4)
	Tortel, Aysen Region	March 20, 2019	SERNAPESCA 2019	Tissue sample in housed in mammals collection Museo Nacional de Historia Natural (MNHN/MAM 1973)
<i>M. grayi</i>	Wreck Point, Posesión Strait of Magellan coast	October, 1979	Sielfeld 1979	Skull and mandible housed in Zoological collection Instituto de la Patagonia Universidad de Magallanes (CE-26)
	Wreck Point, Posesión Strait of Magellan coast	December 12, 1979	Sielfeld 1983	Skull and mandible housed in Zoological collection Instituto de la Patagonia CZIP 0510. Old number = (CE-28). The collect place is was modified (Erika Mutschke, Universidad de Magallanes)
	Catalina Point Strait of Magellan	May 28, 1978	Sielfeld 1983	Skull housed in Zoological collection Instituto de la Patagonia Universidad de Magallanes (CE-35) old number RNP 702
	Guanaqueros Coquimbo Region 30°11'S 71°24'W	December 21, 2018	This work	Specimen housed in Museo Nacional de Historia Natural MNHN/MAM 1972
<i>M. grayi</i> ?	Chanavayita cove, Iquique Region	April 06, 2002	SERNAPESCA 2021 Iquique	Stranded one specimen. Identification by Guillermo Guzmán UNAP
	-29° S	February 1998	Sanino <i>et al.</i> 2007	Sighting, two specimens
<i>M. peruvianus</i>	-20° 26' S, -70° 44' W	December 17, 1997	Sanino <i>et al.</i> 2007	Sighting, one specimen
	-17° 04' S, -71° 23' W	May, 1995	Sanino <i>et al.</i> 2007	Skull
	Pargua bay, Los Lagos Region -41° 45' S, -73° 29' W	March, 1980	Pastene <i>et al.</i> 1990	Stranded one specimen
<i>M. densirostris</i>	Coliumo bay, Concepcion Region 36° -32' S, -73° 57' W	June 19, 1999		Stranded one specimen.
	Vaihu beach Easter island	August 31, 1994	Aguayo <i>et al.</i> 1998	Stranded one specimen. The possible cause of death was a shark attack.
<i>Mesoplodon traversii</i>	Juan Fernández island	June 1986	Val Helden <i>et al.</i> 2002	Specimen housed in Museo Nacional de Historia Natural MNHN/MAM 1156
	-32° 30' S, -88° 42' W	June, 17, 1995	Aguayo <i>et al.</i> 1998	Sighting, one specimen
<i>Mesoplodon</i> sp.	-31° 41' S, -93° 07' W	June, 18, 1995	Aguayo <i>et al.</i> 1998	Sighting, one specimen
	-33° 08' S, -74° 27' W	July, 08, 1995	Aguayo <i>et al.</i> 1998	Sighting, two specimens
	-33° 08' S, 74° 27' W	September, 15, 1995	Aguayo <i>et al.</i> 1998	Sighting, specimen




Figure 3. Bayesian phylogenetic tree resulting from the analysis of cytochrome b gene DNA sequences of specimens of *Mesoplodon*. Numbers at nodes indicate posterior probability (PP; left of the diagonal) and bootstrap support (BS; right of the diagonal) values of the adjacent node. The coloured rectangle indicates the position of the specimen MNHN/MAM 1972 within the phylogenetic tree. The GenBank accession numbers of analyzed sequences are included at terminal labels. The terminal corresponding to the new record of *M. grayi* MNHN/MAM 1972 is highlighted in bold.

Table 2. Measurement of specimen MNHN/MAN 1972 and of other specimens *Mesoplodon grayi* (in cm).

		MNHN/ MAM1972	Female Mean (Thompson <i>et al.</i> 2014)	CE-26 Sielfeld 1979
1	Condylobasal length	45.5	78.2-90.4	73
2	Length of rostrum, tip of beak to line connecting apices of antorbital notches	25.7	53.9-62.9	---
3	Tip of rostrum to most posterior margin of pterygoid near midline	34.0	62.4-74.5	---
4	Tip of rostrum to most posterior extension of wing of pterygoid	35.0	70.6-76.8	---
5	Tip of rostrum to most anterior extension of pterygoid	24.5	45.9-55.7	---
6	Tip of rostrum to most posterior extension of maxillaries between pterygoids on the palate	27.5	51.1-60.7	---
7	Tip of rostrum to most posterior extension of maxillary plate	41.0	73.5-83.85	---
8	Tip of rostrum to anterior margin on superior nares	30.0	59.8-69.6	---
9	Tip of rostrum to most anterior point on premaxillary crest	33.6	63.9-73.2	---
10	Tip of rostrum to most posterior extension of temporal fossae	42.8	79.1-89.5	---
11	Tip of rostrum to most posterior extension of lateral tip of premaxillary crest	33.7	---	---
12	Tip of rostrum to most anterior extension of pterygoid sinus	24.0	49.1-55.7	---
13	Length of temporal fossa	7.8	8.15-10.83	---
14	Length of orbit	7.7	8.8-10.9	8.7
15	Length of right nasal on vertex of skull	4.0	---	---
16	Length of nasal suture	3.8	2.36 - 4.3	---
17	Breadth of skull across postorbital process of frontals	19.8		---
18	Breadth of skull across zygomatic process of squamosals	19.5	28.4-31.8	---
19	Breadth of skull across centres of orbits	18.2		---
20	Least breadth of skull across posterior margins of temporal fossae	16.0		---
21	Greatest span of occipital condyles	---		---
22	Greatest width of an occipital condyle	2.6	3.43-4.53	---
23	Greatest length of an occipital condyle	5.0	1.88-3.36	---
24	Greatest breadth of foramen mangan	---		---
25	Breadht of skull across exoccipitals	16.4	26.6-31.7	---
26	Breadth of nasal vertex	4.1	---	---
27	Least distance between premaxillary crest	1.3	---	---
28	Greatest extension of right premaxillary posterior of right nasal on vertex skull	4.7	---	---
29	Greatest span of premaxillary crest	8.6	12.12-14.66	12.1
30	Least width (strictly transverse) of premaxillae where (and if) they narrow opposite superior nares	8.6	---	---
31	Greatest width of premaxillae anterior to place of measurement N°30	5.3	---	---
32	Width of premaxillae at midlenght of rostrum	1.8	---	---
33	Width of rostrum in apices of antorbital notches	12.7	---	---
34	Width of rostrum in apices of prominential notches (if any)	----	---	---
35	Greatest width of rostrum at midlength of rostrum	2.7	---	---
36	Greatest depth of rostrum at midlength of rostrum	2.7	3.1-3.9	---
37	Greatest transverse width of superior nares	---	---	---
38	Greatest inside width of inferior nares, at apices of pterygoid notches, on the pterygoids	---	---	---
39	Heigth of skull. Distance between vertex of skull and most ventral point on pterygoids	18.2	---	---
40	Greatest width of temporal fossa approximately at right angles to greatest length	5.3	---	---
41	Least distance between (main or anterior) maxillary foramina	2.0	---	---
42	Least distance between premaxillary foramina	1.4	---	---
43	Distance from posterior margin of left maxillary foramen to most anterior extension of left maxillary prominence	6.5	---	---
44	Greatest length of vomer visible at surface of palate	5.3	---	---
45	Amount added to break because of breakage	1.0	---	---
46	Length of tympanic bulla, left	--	---	---
47	Length of tympanic bulla, right	4.9	---	---

Table 2. Continuation

	MNHN/MAM1972
1 Length of mandible	38
2 Greatest length of symphysis	9,6
3 Height of mandible at coronoid process	7,8
4 Outside height of mandible at midlength of alveolus	3,3
5 Inside height of mandible at midlength of alveolus	---
6 Length from most posterior extension of symphysis to most posterior extension of condyle	---
7 Length from posterior margin of alveolus to condyle	---
8 Length of alveolus	3
9 Width of alveolus	0,8
10 Tip of mandible to alveolus	8,1
11 Greatest length of tooth	---
12 Greatest antero-posterior width of tooth at approximately right angles to long axis of tooth	---
13 Greatest breadth of tooth	---

the world (Whitehead et al. 2008; Bernaldo de Quirós et al. 2019; Grove et al. 2020), a pattern that was also observed for cetaceans off the coast of Chile according to records obtained by the Chilean National Fisheries Service (SER-NAPESCA 2021) between 2009 and 2021. There is little information associated with the known Chilean records of the six *Mesoplodon* species that have been recorded. Most of the documented records are based on incomplete bone

material, which prevents the collection of data on coloration, sex and other characteristics. As such, the record presented in this study is relevant as it belongs to Gray's beaked whales. Mann (1958) was one of the first authors to mention the presence of *M. grayi* in Chile. However, Mann (1958) did not provide any geographic references or other information to support his mention. Therefore, we recommend this record to be ignored until further confirmation




Figure 4. World map showing the distribution of *Mesoplodon grayi* records. These are concentrated in three main geographical areas: 1) Australia-Tasmania, which presents the most significant number of records, 2) New Zealand; and finally, 3) the Atlantic coast of South America. Note the records on the Pacific coast of South America are limited to two, one in Peru and the other corresponding to the one presented here. The blue circle indicates the new record of *M. grayi* in Chile (MNHN/MAM 1972); the red circle is the registry in Peru; the yellow circles are the records (strandings and sightings) in other parts of the planet. The black date that indicates the yellow circle in the northern hemisphere, indicates the existence of a specimen deposited in a collection, so it does not represent the distribution of the species

becomes available. There is also a mention for Chanavayita beach (Iquique) which is doubtful and should be taken with caution as the putative record is not associated with any preserved specimen. Therefore, only four records of this species are confirmed for Chile.

Based primarily on coloration, SERNAPESCA assigned specimen MNHN/MAM 1972 to *M. hectori*. The coloration pattern for this species is dark brown, dark grey or black and paler on the ventral surface. The tip of the beak is generally white, unlike the rest of the body (Wilson and Mittermeir 2014). However, specimen MNHN/MAM 1972 presents a different color pattern with a dark colored beak (Figure 2B). The integration of morphological and molecular information allowed us to determine that specimen MNHN/MAM 1972 belongs to *Mesoplodon grayi*. For this species, the body length varies from 474 to 564 cm in adults and 210 to 242 in newborn calf (Reidenberg and Laitman 2009). Specimen MNHN/MAM1972 presents lower values (247 cm). The beak is long and narrow with an almost straight line of the mouth (Redford and Einsenberg 1992; Wilson and Mittermeir 2014), unlike specimen MNHN/MAM 1972 which has a short beak and a curved buccal line, characteristics that could be attributed to its juvenile condition. The literature reports small teeth at the maxillary level (Von Haast 1876; Sielfeld 1979, 1983; Reyes and Molina 1997), which are also observable in the reported specimen (Figure 2). The skull shows the anterior border of the pterygoid sinus at the level of the maxillary foramen (= dorsal infraorbital foramen according to Mead and Fordyce 2009). The left premaxillary ridge is smaller and less marked than the right one. Finally, mature males have a well-developed pair of teeth in the jaw, while in females these teeth are smaller and not always visible as are covered by the gum (Mead 1989); the latter is the case of MNHN/MAM 1972. The difference in the location of the teeth in the jaw between *M. hectori* and *M. grayi* is a distinctive character. In *M. hectori*, the teeth are located at the tip of the mandible (apical), while in *M. grayi*, they are immediately anterior to the mandibular symphysis (Reyes and Molina 1997). It is important to mention that position of the teeth in *M. grayi* varies (Von Haast 1876; Robson 1975; Sielfeld 1979; Reyes 1990). The difference observed by these authors may be due to ontogenetic differences of the species. It should also be considered that there are other sources of phenotypic variation that could explain the distinction of specimen MNHN/MAM 1972. Thompson et al. (2014) mentioned the existence of sexual dimorphism expressed in the skull for this species in two areas of New Zealand, as well as geographic variation. The differences found in specimen MNHN/MAM 1972 may be due to its juvenile condition (Table 2).

Worldwide, 158 specimens of *Mesoplodon grayi* are currently available in collections (Figure 4). These specimens mainly come from three well-defined geographic areas: 1) Australia-Tasmania, which has the highest number of preserved and observed specimens, 2) New Zealand, and 3) the Atlantic coast of South America (GBIF 2019). Only one

record is known, from the coast of Peru (Paracas), from the Pacific coast of South America; corresponds to a stranded female of *M. grayi* identified by external phenotype and skull measurements (Reyes 1990). As such, records on the Pacific coast of South America are scarcely reaching to six with this new Chilean record for *M. grayi* which is also very valuable as it is the first of a complete specimen from which molecular data are presented.

Acknowledgments

Our gratitude to P. Schiappacasse for the support in the transfer of the specimen from Coquimbo and to P. Rivera for the information provided on the specimen from Iquique. Our gratitude to E. Mutschke, from the Institute of Patagonia (ISP), the University of Magallanes for providing us with photographs of material from the ISP collection as well as important data. A special thanks to V. Rivas who made and edited the photographs of the MNHN/MAM 1972 skull. Molecular analyses of the specimen were made possible thanks to the support of the FONDECYT 1180366 (GD) project.

Literature cited

- AGUAYO, A. 1975. Progress report on small cetaceans research in Chile. *Journal Fisheries Research Board of Canada* 32:1123-1143.
- AGUAYO, A., ET AL. 1998. Observaciones de cetáceos realizadas entre Valparaíso e Isla de Pascua, Chile, durante los inviernos de 1993, 1994 y 1995. *Revista de Biología Marina y Oceanografía* 33:101-123.
- AKAIKE, H. 1973. Information theory and an extension of the maximum likelihood principle. Pp. 267-281, in *Proceedings of the 2nd International Symposium on Information Theory* (Petrov, B. N., and F. Csaki, eds.). Akademiai Kiado. Budapest, Hungary.
- BAKER, A. N., AND A. L. VAN HELDEN. 1999. New record of Beaked whales, genus *Mesoplodon*, from New Zealand (Cetacea: Ziphiidae). *Journal of the Royal Society of New Zealand* 29:235-244.
- BERNALDO DE QUIRÓS, Y., ET AL. 2019. Advances in research on the impacts of anti-submarine sonar on beaked whales. *Proceedings Royal Society B*. 286:1-9
- BERTA, A., J. L. SUMICH, AND K. M. KOVACS. 2015. *Marine Mammals: evolutionary biology*. Academic Press Elsevier. London, U.K.
- BESHARSE, J. C. 1971. Maturity and sexual dimorphism in the skull, mandible, and teeth of the Beaked whale, *Mesoplodon densirostris*. *Journal of Mammalogy* Vol. 52:297-315.
- CANTO, J., P. RUIZ, AND J. C. CÁRDENAS. 1991. Necropsia de Ballena Franca Austral (*Eubalaena australis*) y consideraciones sobre manejo de la especie. *Boletín del Museo Nacional de Historia Natural, Chile* 42:105-111.
- CANTO, J., P. RUIZ, AND J. YÁÑEZ. 1992. Registros de nuevas especies de cetáceos para la costa de Chile y antecedentes del grupo. *Boletín del Museo Nacional de Historia Natural, Chile* 43:105-115.
- CANTO, J., AND J. YÁÑEZ. 2009. Cetáceos. Pp. 159-192, in *Mamíferos de Chile* (Pedreros, A. H. M., and J. Y. Valenzuela, Eds.). Editorial CEA. Valdivia, Chile.
- CAPPOZZO, H. L., ET AL. 2005. Biological data on two Hector's Beaked whales, *Mesoplodon hectori*, stranded in Buenos Aires, province, Argentina. *Latin American Journal of Aquatic Mammals* 4:113-128

- CÁRDENAS, J. C., M. E. STUTZIN, J. E. OPORTO, C. CABELLO AND D. TORRES. 1986. Manual de identificación de los cetáceos chilenos. Proyecto WH-445 Cetáceos-Chile, World Wildlife Fund-US – CODEFF. Available at: correo@electronico
- COX, T.M., ET AL. 2006. Understanding the impacts of anthropogenic sound on beaked whales. *Journal of Cetacean Research and Management* 7:177-187.
- DALEBOUT, M. L., A. L. VAN HELDEN, K. VAN WAEREBEEK AND A. N. BAKER. 1988. Molecular genetic identification of the Southern hemisphere Beaked whales (Cetacea: Ziphiidae). *Molecular Ecology* 7:687-694.
- DALEBOUT, M. L., J. G. MEAD, C. S. BAKER, A. N. BAKER AND A. L. VAN HELDEN. 2002. A new species of Beaked whale *Mesoplodon perrini* sp.n. (Cetacea: Ziphiidae) discovered through phylogenetic analyses of mitochondrial DNA sequences. *Marine Mammal Science* 18:577-608.
- DALEBOUT, M. L., C. S. BAKER, J. G. MEAD, V. G. COCKCROFT, AND T. K. YAMADA. 2004. A comprehensive and validated molecular taxonomy of beaked whales, family Ziphiidae. *Journal of Heredity* 95:459-473.
- DALEBOUT, M. L., ET AL. 2014. The resurrection of *Mesoplodon hotaula* Deraniyagala 1963: A new species of beaked whale in the tropical Indo-Pacific. *Marine Mammals Science* 30:1081-1108.
- DEL CASTILLO, D. 2016. Desarrollo ontogenético craneano en cetáceos odontocetos del hemisferio sur. Tesis Doctoral. Universidad de Buenos Aires. Facultad de Ciencias Exactas y Naturales.
- D'ELIA, G., ET AL. 2020. Lista actualizada de los mamíferos vivientes de Chile. *Boletín del Museo Nacional de Historia Natural, Chile* 69:67-98.
- EVANS, K., ET AL. 2005. Periodic variability in cetacean strandings: Links to large-scale climate events. *Biology Letters* 1:147-150.
- GBIF. 2019. GBIF.org (October 09 2019) GBIF Occurrence Download <https://doi.org/10.15468/dl.nbn05j>.
- GROVE, T., C. SENGLAT, M. PETITGUYOT, D. KOSIBA, AND M. H. RASMUSSEN. 2020. Mass stranding and unusual sightings of northern bottlenose whales (*Hyperoodon ampullatus*) in Skjálfandi Bay, Iceland. *Marine Mammal Science* 36:1033-10411.
- HELDEN, A. L., ET AL. 2002. Resurrection of *Mesoplodon traversii* (Gray, 1874), senior synonym of *M. bahamondi* Reyes, Van Waerebeek, Cárdenas and Yáñez, 1995 (Cetacea: Ziphiidae). *Marine Mammal Science* 18:609-621.
- KATOH, K., AND D. M. STANDLEY. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* 30:772-780.
- KATOH, K., AND H. TOH. 2008. Improved accuracy of multiple ncRNA alignment by incorporating structural information into a MAFFT-based framework. *BMC Bioinformatics* 9:212.
- LAPORTA, P., R. PRADERI, V. LITTLE, AND A. LE BAS. 2005. An Andre's Beaked whale *Mesoplodon bowdoini* (Cetacea, Ziphiidae) stranded on the Atlantic coast of Uruguay. *Latin American Journal of Aquatic Mammals* 4:101-111.
- LEENEY, R. H., ET AL. 2008. Spatio-temporal analysis of cetacean strandings and bycatch in a UK fisheries hotspot. *Biodiversity Conservation* 17:2323-2338.
- LITCHER, A. 1986. Records of Beaked whales (Ziphiidae) from the western South Atlantic. *Scientific Report of the Whales Research Institute* 37:109-127.
- MANN, G. 1958. Clave de determinación para las especies de Mamíferos Silvestres de Chile. *Investigaciones. Zoológicas Chilenas, Santiago* 3:38.
- MACLEOD, C. D. 2017. Beaked Whales, Overview. Pp. 80-88, in *Encyclopedia of Marine Mammals*. Academic Press. San Diego. U.S.A.
- MCLEOD, C. D. 2009. Global climate change, range changes and potential implications for the conservation of marine cetaceans: a review and synthesis. *Endangered Species Research* 7:125-136.
- MCLEOD, C. D., ET AL. 2006. Known and inferred distributions of Beaked whale's species (Cetacea: Ziphiidae). *Journal Cetacean Research Management* 7:271-286.
- MEAD, J. G., AND R. E. FORDYCE. 2009. The therian skull: a lexicon with emphasis on the odontocetes. *Smithsonian Contributions to Zoology*. Scholarly Press <https://doi.org/10.5479/si.00810282.627>
- MEAD, J. G. 1989. Beaked whales of the genus *Mesoplodon*. Pp. 349-430, in *Handbook of Marine Mammals*. (Ridgway, H., and R. Harrison, Eds.). Vol. 4. Academic Press. New York, U.S.A.
- MEAD, J. G., AND A. N. BAKER. 1987. Notes on the rare Beaked whale, *Mesoplodon hectori* (Gray). *Journal of the Royal Society of New Zeland* 17:303-312.
- MOORE, K. M., C. A. SIMEONE, AND R. L. BROWNELL JR. 2018. Strandings. Pp. 945-951, in *Encyclopedia of Marine Mammals*. (Würsig, B., J. G. M. Thewissen, and K. M. Kovacs, eds.). Academic Press, London, United Kingdom.
- NOBUYUKI, M., I. NAKAMURA, S. TANABE, AND R. TATSUKAWA. 1987. A stranding of *Mesoplodon stejnegeri* in the Maizuru bay, Sea of Japan. *Scientific Report of the Whales Research Institute* 37:91-105.
- NORMAN, S., AND J. G. MEAD. 2001. *Mesoplodon europaeus*. *Mammalian Species* 668:1-5.
- PASTENE, L. A., K. NUMACHI, M. JOFRE, M. ACEVEDO AND G. JOYCE. 1990. First record of the Blainville's Beaked Whale, *Mesoplodon densirostris* Blainville, 1817 (Cetacea, Ziphiidae) in the eastern south pacific. *Marine Mammal Science* 6:82-84.
- PIKESLEY, S. K., ET AL. 2011. Cetacean sightings and strandings: evidence for spatial and temporal trends? *Journal of the Marine Biological Association UK* 92:1809-1820.
- PITMAN, R. 2009. *Mesoplodon* Beaked Whales. Pp. 595-602, in *Encyclopedia of Marine Mammals*. (Würsig, B., J. G. M. Thewissen, and K. M. Kovacs, Eds.). Academic Press. London, U.K..
- REDFORD, K. H., AND J. F. EISENBERG. 1992. *Mammals of the Neotropics. The outhern Cone Volume 2. Chile, Argentina, Uruguay, Paraguay*. The University of Chicago Press. Chicago, U.S.A.
- REIDENBERG, J. S. and J. T. LAITMAN. 2009. Cetacean prenatal development. Pp. 220-230, in *Encyclopedia of Marine Mammals*. (Würsig, B., J. G. M. Thewissen, and K. M. Kovacs, eds.). Academic Press, London, U. K.
- ROBSON, F. D. 1975. On Vestigial and normal teeth in the scamper-down beaked whale, *Mesoplodon Ggayi*. *Tuatara* 21:105-107.
- RONQUIST, F. and J. P. HUELSENBECK. 2013. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19:1572-1574.
- REYES, J. C. 1990. Gray's beaked whale *Mesoplodon grayi* in the South East Pacific. *Mammalian Biology (früher Zeitschrift für Säugetierkunde)* 55:139-141.

- REYES, J. C., AND D. MOLINA. 1997. Clave artificial para la identificación de cráneos de cetáceos del Pacífico sureste. *Boletín del Museo Nacional de Historia Natural, Chile* 46:95-119.
- REYES, J. C., K. VAN WAEREBEEK, J. C. CÁRDENAS, AND J. L. YÁÑEZ. 1995. *Mesoplodon bahamondi* sp. n. (Cetacea: Ziphiidae), a new living beaked whale from Juan Fernández Archipelago, Chile. *Boletín del Museo Nacional de Historia Natural, Chile* 45:31-44.
- REYES, J. C., AND K. VAN WAEREBEEK. 2018. The lesser Beaked whale *Mesoplodon peruvianus* Reyes, Mead & Van Waerebeek 1991 revisited, with biological observation on new specimens from Peru. *Journal of Marine Biology & Oceanography* 7:1-10.
- ROSS, G. J. B. 1984. The smaller cetaceans of the south east coast of southern Africa. *Annals of the Cape Provincial Museums, Natural History* 15:173-410.
- ROSS, G. J. B. 1970. The occurrence of Hector's Beaked whale *Mesoplodon hectori* (Gray) in South African waters. *Annals of the Cape Provincial Museum (Natural History)* 8:195-204.
- SANINO, G. P, J. YÁÑEZ, AND K. VAN WAEREBEEK. 2007. A first confirmed specimen record in Chile, and sightings attributed to the lesser beaked whale *Mesoplodon peruvianus* Reyes, Mead and Van Waerebeek, 1991. *Boletín del Museo Nacional de Historia Natural, Chile* 56:89-96.
- SERNAPESCA. 2021. Base de datos 2009-2021. Registro de varamientos. Web: <http://www.sernapesca.cl/informacion-utilidad/registro-de-varamientos>.
- SIELFELD, W. 1979. Consideraciones acerca de tres especies de *Mesoplodon* Gervais (Cetacea: Ziphiidae) presentes en aguas chilenas. *Anales del Instituto de la Patagonia* 10:181-187.
- SIELFELD, W. 1980. Mamíferos marinos en las colecciones y museos de Chile. *Anales del Instituto de la Patagonia* 11:273-279.
- SIELFELD, W. 1983. Mamíferos marinos de Chile. Ediciones de la Universidad de Chile.. Santiago, Chile.
- THOMPSON, K. F., K. RUGGIERO, C. D. MILLAR, R. CONSTANTINE, AND A. L. VAN HELDEN. 2014. Large-scale multivariate analysis reveals sexual dimorphism and geographic differences in the Gray's beaked whale. *Journal of Zoology* 294:13-21.
- Trifinopoulos, J., L. T. Nguyen, A. von Haeseler, and B. Q. Minh. 2016. W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Research* 8:232-235
- VAN HELDEN, A. L., ET AL. 2002. Resurrection of *Mesoplodon traversii* (Gray, 1874), senior synonym of *M. bahamondi* Reyes, Van Waerebeek, Cardenas and Yanez, 1995 (Cetacea: Ziphiidae). *Marine Mammal Science* 18:609-621.
- VENEGAS, C., AND W. SIELFELD. 1978. Registros de *Mesoplodon layardii* y otros cetáceos en Magallanes. *Anales Instituto Patagonia (Punta Arenas, Chile)* 9:171-177.
- VON HAAS, J. 1876. On a new Ziphioid whale. *Proceedings of the Zoological Society of London* 4:7-13.
- WHITEHEAD, H., B. MCGILL, AND B. WORM. 2008. Cetacean diversity of deep-water cetaceans in relation to temperature: implications for ocean warming. *Ecology Letters* 11:1198-1207.
- WILSON, D. E., AND R. A. METTERMEIER. 2014. Handbook of the Mammals of the World Vol 4 Sea Mammals. Lynx Edicions. Barcelona. Spain.
- YAMADA, T.K., ET AL. 2019. Description of a new beaked whale (*Berardius*) found in the North Pacific. *Scientific Reports* 9:1-14.
- ZERBINI, A. N., AND E. SECCHI. 2001. Occurrence of Hector's beaked whale, *Mesoplodon hectori* in southern Brazil. *Aquatic Mammals* 27:149-153.

Associated editor: Juan Pablo Gallo-Reymoso

Submitted: January 5, 2022; Reviewed: February 26, 2022

Accepted: July 17, 2022; Published on line: August 11, 2022